

Feed More Milk without Scours

Cold weather arrives. You decide to feed more milk/milk replacer. Soon after making the change your treatable scours rate goes up too much to be acceptable.

What are the differences among farms that have this problem and those that feed milk/milk replacer at higher volumes without diarrhea issues among young calves?

The most common differences

Low Scours Rate	High Scours Rate
1. Milks fresh cows as soon as possible after calving, nearly all of them within six hours post-calving.	1. Milks fresh cows next regularly scheduled milking.
2. Checks colostrum quality and uses highest quality for first feeding.	2. Does not check colostrum quality.
3. Feeds colostrum as soon as possible after birth, always within first four hours.	3. Feeds colostrum at next regular calf feeding time.
4. Feed 3.5-4 quarts colostrum (large breeds)	4. Feeds 1.5-2 quarts of colostrum.
5. Checks colostrum cleanliness with regular culturing.	5. Does not check colostrum for bacteria content.
6. Checks for successful passive transfer of immunity on a regular basis.	6. Checks for successful passive transfer of immunity only if there is a "problem."
7. Cleans colostrum and milk handling equipment after every use following an accepted cleaning protocol that is written and posted.	7. Cleans colostrum and milk handling equipment as convenient with no regular protocol.
8. Checks milk or milk replacer cleanliness with regular culturing.	8. Does not check milk or milk replacer for bacteria content.
9. Feeds preweaned calves enough milk or milk replacer to support at least 1 pound a day gain all seasons of the year.	9. Feeds preweaned calves milk or milk replacer at a rate such that calves do not gain weight some seasons of the year.
10. Keeps calf housing clean.	10. Houses calves in a high bacteria environment.

How Realistic is it to try Feeding at a Higher Volume?

Following all the practices in the left-hand column above does not guarantee that none of your calves will have scours. In contrast, the chances for scours do go up as your practices look more and more like the ones in the right-hand column.

Feeding calves is always like walking a tight-rope. You are trying to maintain a balance. As you increase milk or milk replacer feeding volumes the chances of losing your balance go up. That is, the calves have diarrhea. This requires better management skills.

Key Skills:

- Be able to feed different volumes of milk to calves – not every calf receives the same amount. While there are a few exceptions most calf feeding programs that feed more than the traditional two quarts twice daily increase volume as calves grow. Lots of folks mark individual or groups of pens to receive a specific amount per feeding.
- Be able to feed consistent volumes of milk. This means delivering each feeding within one cup of the intended volume. For example, when feeding three quarts at one feeding the actual amount delivered does not vary more than 2.75 to 3.25 quarts.
- Be able to deliver milk replacer mixed at the same concentration at every feeding. A significant step in achieving this consistency is having an accurate set of scales that are used all the time to measure milk replacer powder.
- Be able to deliver milk or milk replacer at the same temperature at every feeding. My goal is to achieve delivery temperatures in the range of 100-105 F. In cold weather conditions this may mean delivering liquid feeds in multiple batches.
- Be able to observe and diagnose scours in calves. Prompt diagnosis and treatment is always important. Equally important is watching a group of calves the first few days after their ration has been bumped up in volume.

Many folks have observed that it is a good practice to temporarily drop back volume fed for a few days when a calf scours after a ration increase. My personal experience suggests that at least one out of twenty calves will experience what is often called “nutritional” scouring even when volume increases are as small as 0.5 quart per feeding.

Sam Leadley, Calf & Heifer Management Specialist

sleadley@yahoo.com www.calffacts.com

© Attica Vet. Assoc. 2013 All Rights Reserved.